



c09-c-302

3218

**BOARD DIPLOMA EXAMINATION, (C-09)
OCT/NOV—2013
DCE—THIRD SEMESTER EXAMINATION
ENGINEERING MATHEMATICS—II**

Time : 3 hours]

[Total Marks : 80

PART—A

- Instructions :** (1) Answer **all** questions.
(2) Each question carries **three** marks.
(3) Answers should be brief and straight to the point and shall not exceed *five* simple sentences.

1. Evaluate :

$$\frac{\tan^{-1} x}{1-x^2} dx$$

2. Evaluate :

$$\sqrt{1-\cos 2x} dx$$

3. Evaluate :

$$\sec(4x-7) \tan(4x-7) dx$$

*

C09-C-302

4. Evaluate :

$$\frac{dx}{\sqrt{3-x^2}}$$

5. Evaluate :

$$x^5 - \frac{4}{x} + 3 \cos x \, dx$$

6. Evaluate :

$$\int_0^1 (x^4 - 1) dx$$

7. Write the formula to find the RMS value of $y = f(x)$ between $x = a$ and $x = b$.

8. Solve :

$$x^5 dy - y^5 dx = 0$$

9. Find the particular integral of $(D^2 - 4)y = \sin 2x$.

10. Form the differential equation of the family of curves $y = Ae^{2x} + Be^{-2x}$, where A, B are arbitrary constants.

*

PART—B

- Instructions :** (1) Answer *any five* questions.
(2) Each question carries **ten** marks.
(3) Answers should be comprehensive and the criterion for valuation is the content but not the length of the answer.

11. (a) Evaluate :

$$\int \sin^2 x \cos^3 x \, dx$$

(b) Evaluate :

$$\int \sin 5x \cos 2x \, dx$$

12. (a) Evaluate :

$$\int \frac{1}{x^2 + 6x + 13} \, dx$$

(b) Evaluate :

$$\int x^3 e^x \, dx$$

13. (a) Find the volume of the solid obtained by revolving the ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

about x -axis.

(b) Find the RMS value of $\sqrt{\log x}$ over the range $x = 1$ to $x = e$.

14. (a) Evaluate :

$$\int_0^{\pi/2} \frac{\sin^{12} x}{\sin^{12} x + \cos^{12} x} \, dx$$

(b) Using the method of integration, find the area bounded by the circle $x^2 + y^2 = a^2$.

*

15. (a) Solve :

$$(D^2 - 1)y = \cos x$$

(b) Solve :

$$(D^2 - 3D - 2)y = x^2$$

16. (a) Solve :

$$\frac{dy}{dx} - \frac{2y}{x} = 3x$$

(b) Solve :

$$(4D^2 - 4D - 3)y = e^{2x}$$

17. Solve :

$$x^2 y dx - (x^3 - y^3) dy$$

18. (a) Use Simpson's rule to find the value of

$$\int_1^2 \frac{1}{x^2} dx$$

by taking five ordinates.

(b) Solve :

$$e^y dx - (xe^y - 2y) dy = 0$$
